

University of Groningen

Insulin analogues use in pregnancy among women with pregestational diabetes mellitus and risk of congenital anomaly

Wang, Hao; Wender-Ozegowska, Ewa; Garne, Ester; Morgan, Margery; Loane, Maria; Morris, Joan K; Bakker, Marian K; Gatt, Miriam; de Walle, Hermien; Jordan, Susan

Published in:
BMJ Open

DOI:
[10.1136/bmjopen-2016-014972](https://doi.org/10.1136/bmjopen-2016-014972)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wang, H., Wender-Ozegowska, E., Garne, E., Morgan, M., Loane, M., Morris, J. K., Bakker, M. K., Gatt, M., de Walle, H., Jordan, S., Materna-Kiryluk, A., Nelen, V., Thys, G., Wiesel, A., Dolk, H., & de Jong-van den Berg, L. T. W. (2018). Insulin analogues use in pregnancy among women with pregestational diabetes mellitus and risk of congenital anomaly: a retrospective population-based cohort study. *BMJ Open*, 8(2), [014972]. <https://doi.org/10.1136/bmjopen-2016-014972>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

BMJ Open Insulin analogues use in pregnancy among women with pregestational diabetes mellitus and risk of congenital anomaly: a retrospective population-based cohort study

Hao Wang,¹ Ewa Wender-Ozegowska,² Ester Garne,³ Margery Morgan,⁴ Maria Loane,⁵ Joan K Morris,⁶ Marian K Bakker,⁷ Miriam Gatt,⁸ Hermien de Walle,⁷ Susan Jordan,⁹ Anna Materna-Kiryluk,¹⁰ Vera Nelen,¹¹ Guy Thys,¹¹ Awi Wiesel,¹² Helen Dolk,¹³ Lolkje T W de Jong-van den Berg¹

To cite: Wang H, Wender-Ozegowska E, Garne E, *et al.* Insulin analogues use in pregnancy among women with pregestational diabetes mellitus and risk of congenital anomaly: a retrospective population-based cohort study. *BMJ Open* 2018;**8**:e014972. doi:10.1136/bmjopen-2016-014972

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2016-014972>).

Received 6 December 2016
Revised 8 January 2018
Accepted 9 January 2018

ABSTRACT

Objectives To evaluate the risk of major congenital anomaly associated with first-trimester exposure to insulin analogues compared with human insulin in offspring of women with pregestational diabetes.

Design and setting A population-based cohort of women with pregestational diabetes (n=1661) who delivered between 1996 and 2012 was established retrospectively from seven European regions covered by the European Surveillance of Congenital Anomalies (EUROCAT) congenital anomaly registries.

Primary outcome measures The risk of non-chromosomal major congenital anomaly in live births, fetal deaths and terminations for a fetal anomaly exposed to insulin analogues in the first trimester of pregnancy was compared with the risk in those exposed to human insulin only.

Results During the first trimester, 870 fetuses (52.4%) were exposed to human insulin only, 397 fetuses (23.9%) to insulin analogues only and 394 fetuses (23.7%) to both human insulin and insulin analogues. The risk of major congenital anomaly in fetuses exposed to insulin analogues only was lower than those exposed to human insulin only; the relative risk adjusted for glycaemic control and region was 0.56 (95% CI 0.29 to 1.06). The significantly lower risk related to exposure of insulin analogues only was observed in congenital heart defects: adjusted relative risk 0.14 (95% CI 0.03 to 0.62).

Conclusions In this retrospective population-based cohort study across Europe, first-trimester exposure to insulin analogues did not increase the risk of major congenital anomaly compared with exposure to human insulin. A possible lower risk of congenital heart defects among fetuses exposed to insulin analogues only deserves further investigation.

Strengths and limitations of this study

- A large retrospective population-based cohort of diabetic pregnancies from multiple centres that represent geographically distinct areas where patterns of diabetic treatment are different.
- The valid information on congenital anomaly was obtained by linkage to the EUROCAT database and by reclassification the cases with written text descriptions of a congenital anomaly based on EUROCAT subgroup definition.
- The data sources which were based on medical records and healthcare databases had incomplete data on haemoglobin A1c (HbA1c) and pregnancy planning, but the association of congenital anomalies and insulin analogues was examined only among women who had had a HbA1c in the first trimester.
- The database does not contain information on lifestyle factors (eg, smoking, obesity, alcohol), preconception care (eg, screening and treatment of complications of diabetes, folic acid supplementation) which are also known to lower the risk of congenital anomaly.

including stillbirth, major congenital anomalies, and neonatal morbidity and mortality^{1–5} compared with women without diabetes. Isolated congenital heart defects account for one-third of all congenital anomalies in pregnancies with diabetes and a quarter in pregnancies without diabetes.⁶ Although the mechanisms underlying the associations of DM with congenital anomalies are not completely understood, there is evidence for a positive association between hyperglycaemia during embryogenesis and the risk for congenital anomalies.^{7,8} Preconception care focusing on glycaemic control can improve pregnancy outcome and reduce



For numbered affiliations see end of article.

Correspondence to

Dr Hao Wang; hao.wang@rug.nl

INTRODUCTION

Women with pregestational diabetes mellitus (DM) (type 1 or type 2) have a higher risk of serious adverse pregnancy outcomes,

the frequency of congenital anomalies in women with DM.⁹

Clinical and experimental studies report that insulin analogues result in improved glycaemic control, more stable glycaemia, less glucose spikes, fewer hypoglycaemic episodes and improved patient satisfaction compared with the use of human insulin,^{10–12} which might have benefits for pregnant women. However, studies on exposure to insulin analogues during pregnancy are small and underpowered to evaluate the risk of specific major congenital anomalies.¹³ We conducted a study using a cohort of pregnancies with diabetes across Europe to evaluate the risk of major congenital anomalies associated with insulin analogue use in the first trimester of pregnancy compared with the use of human insulin.

METHODS

Study population

A multicentre, retrospective cohort study was performed in seven regions in Europe, belonging to the EUROCAT network of population-based congenital anomalies registries, all obliged to value the same definitions, standards and procedures. Inclusion criteria were women with pregestational diabetes who had been referred to the hospitals in the region of study centre and delivered between 1996 and 2012. The study included live births, fetal deaths (spontaneous abortion, stillbirths) and terminations for a fetal anomaly. Selection of study centres that captured pregnant women with DM for this study was on the basis of contacts within the EUROCAT network. The population of the study centre had to be covered by the EUROCAT network in order to evaluate the pregnancy outcome in terms of congenital anomaly. Each centre should include information on the key variables for evaluating insulin analogues use in the first trimester. Information on maternal drug exposure and other confounders was obtained from medical records. A total of 1877 fetuses of women with pregestational diabetes were enrolled in the study. We excluded 160 fetuses exposed to oral glucose-lowering medications in the first trimester and 56 fetuses with unspecified maternal treatment of diabetes. The final cohort for the study included 1661 fetuses from 1634 pregnancies.

Database of cohort with diabetes

Data on maternal demographics, type of diabetes, treatment, that is, the name of the medication or complete seven-digit Anatomical Therapeutic Chemical classification code, including the dispensing date and number of days of supply, duration of diabetes before conception, planned pregnancy, haemoglobin A1c (HbA1c) level and neonatal outcomes were collected from local medical records.

EUROCAT database

Details of the EUROCAT central database have been described previously.¹⁴ Congenital anomalies among

live births, fetal deaths and terminations for a fetal anomaly are standardly recorded according to EUROCAT Guide 1.3.¹⁵ One syndrome and up to eight malformations are coded by International Classification of Diseases, Ninth Revision (ICD9) or 10th Revision (ICD10) codes. Minor anomalies, according to the EUROCAT classification system, are excluded. The follow-up period for inclusion of congenital anomaly in the EUROCAT varies among study regions: up to 1 week after birth in Mainz, up to 1 year after birth in Antwerp, Malta and Wales, up to 2 years after birth in Poznan, up to 5 years after birth in Funen and up to 16 years after birth in Northern Netherlands.

Exposure definition

The relevant exposure period was defined from the date of the last menstrual period to week 13 of pregnancy (first trimester). All fetuses were classified according to first-trimester maternal treatment of diabetes. The following three groups were included in the analysis: (1) fetuses exposed to human insulin only, (2) fetuses exposed to insulin analogue only and (3) fetuses exposed to both human insulin and insulin analogue. The reference group was fetuses exposed to human insulin only in the first trimester.

Outcome definitions

The primary outcome of interest was defined as the diagnosis of a major congenital anomaly in the infant/fetus, while the secondary outcome was a diagnosis of congenital heart defects. Written descriptions of congenital anomalies based on medical records for live births, fetal deaths and terminations for a fetal anomaly at any gestational age were recorded in the database of cohort with diabetes. A two-stage process was used to identify and confirm fetuses with a major congenital anomalies. First, major congenital anomalies were identified by linking the database of cohort with diabetes to the EUROCAT central database. Second, unlinked cases with a written description of congenital anomalies were reviewed by a paediatrician (EG), blind to exposure status, to classify them into major versus minor congenital anomalies. All major congenital anomalies were grouped according to the EUROCAT classification system.¹⁵ Fetuses with minor anomalies according to EUROCAT criteria were counted as not having a congenital anomaly diagnosis. Chromosomal anomalies (n=5) were excluded from major congenital anomalies and congenital heart defects in the analysis of risk estimation and EUROCAT organ subgroup categorisation.

Preterm birth was defined as delivery before 37 weeks of gestation. We determined the percentile for birth weight for each neonate based on the gestational age using Dutch sex-specific standards.¹⁶ Large for gestational age (LGA) was defined as those above the 90th percentile weight by gestational age and sex. An adverse pregnancy outcome was defined as a major congenital anomaly or a fetal death, which included spontaneous abortion (gestational age <20 weeks) and stillbirth (gestational age ≥20 weeks).



Statistical analyses

We compared the distributions of maternal and neonatal characteristics among the three exposure groups. One-way analysis of variance was used to compare the mean differences in maternal age at delivery, gestational age and birth weight for live births only between exposure groups, and the Scheffe multiple comparison test was used for comparison with the reference group (human insulin only). Multinomial logistic regression was used to identify the differences in categorical variables between exposure groups. Associations between congenital anomalies/congenital heart defects and treatment of diabetes were estimated by calculating both crude and adjusted ORs and their 95% CIs using logistic regression models. ORs were adjusted for study centre (using a random-effect model) and HbA1c levels in the first trimester. HbA1c levels were collected from 3 months before pregnancy until delivery and a mean value was calculated for each trimester. The mean HbA1c level in the first trimester was classified into four categories: mean HbA1c ≤ 43 mmol/mol ($\leq 6.1\%$), mean HbA1c 44–53 mmol/mol (6.2%–7.0%), mean HbA1c 54–67 mmol/mol (7.1%–8.3%) and mean HbA1c > 67 mmol/mol ($> 8.3\%$) and unknown. Missing HbA1c levels in the first trimester (n=614) were excluded from the main analysis of risk estimation of major congenital anomalies/congenital heart defects, because the first-trimester level of HbA1c is essential when considering congenital abnormalities. A sensitivity analysis was conducted by including missing HbA1c as a separate group in the multivariate logistic model (online supplementary table). Analyses were performed using STATA V.13.

RESULTS

During the first trimester, 870 (52.4%) fetuses were exposed to human insulin only, 397 (23.9%) fetuses to insulin analogues only, 394 (23.7%) fetuses to both human insulin and insulin analogues. Of 397 fetuses exposed to insulin analogues only, 277 were exposed to short-acting insulin analogues only: 169 to insulin lispro, 106 to insulin aspart and 1 to insulin glulisine. Of 120

fetuses exposed to short-acting insulin analogues with long-acting insulin analogues, 93 were exposed to insulin glargine and 27 exposed to insulin detemir. The proportion of insulin analogues only exposed fetuses increased from 3.2% in 1996–2002 to 20.8% in 2003–2007, and to 39.3% in the period 2008–2012. We found regional differences in the proportion of exposure to insulin analogues only in the first trimester ranging from 3.8% in Malta to 59.6% in Wales (table 1).

Compared with fetuses exposed to human insulin only in the first trimester, fetuses exposed to insulin analogues only were more common among women who planned their pregnancies (49.4% vs 23.9%, $P<0.0001$), with type 1 diabetes (95.5% vs 90.9%, $P<0.004$) and who had a longer duration of diabetes (64.2% vs 46.6%, $P<0.0001$). Overall, the mean HbA1c value in the first trimester was 55.5 mmol/mol (7.2%) (SD 16.1). Fetuses exposed to insulin analogue only in the first trimester were more likely to be in mean HbA1c 44–53 mmol/mol (6.2%–7.0%) group compared with those exposed to human insulin only (29.5% vs 13.8%, $P<0.001$) (table 2). Neonatal outcomes in terms of gender, LGA, death in the first week were similar among all groups. The proportion of spontaneous abortion was 1.5% (n=6) among fetuses exposed to insulin analogues only, and 1.3% (n=5) among fetuses exposed to both insulin analogues and human insulin. These rates were significantly higher than fetuses exposed to human insulin only 0.1% (n=1) (table 2). Major congenital anomalies were observed in 70 (8.0%) fetuses with human insulin only exposure, 15 (3.8%) fetuses with insulin analogue only exposure and 19 (4.8%) fetuses with human insulin and insulin analogue exposure. Fetuses from Northern Netherlands had the lowest rate of congenital anomaly overall (2.1%, 95% CI 0.9% to 4.7%). The highest rate of congenital anomaly was seen in Antwerp (12.3%, 95% CI 7.3% to 19.8%).

A total of 1047 fetuses whose mother had a first-trimester HbA1c were included in the analysis of the risk estimation of congenital anomalies/congenital heart defects. Compared with fetuses exposed to human insulin only, those exposed to insulin analogues only had a lower

Table 1 Diabetic treatment in the first trimester by centre

Centre	Birth year	Human insulin only, N=870		Insulin analogue only, N=397		Human insulin and insulin analogue, N=394	
		n	%	n	%	n	%
Funen (N=206)	2000–2011	150	72.8	13	6.3	43	20.9
Northern Netherlands (N=270)	1999–2012	63	23.3	145	53.7	62	23.0
Malta (N=132)	1999–2011	127	96.2	5	3.8	0	0
Antwerp (N=99)	1997–2011	31	31.3	39	39.4	29	29.3
Mainz (N=204)	1996–2012	154	75.5	32	15.7	18	8.8
Wales (N=146)	1996–2012	54	37.0	87	59.6	5	3.4
Poznan (N=606)	1999–2012	292	48.2	77	12.7	237	39.1

Table 2 Maternal and infant characteristics by treatment of diabetes in the first trimester†

	Human insulin only, n (%)	Insulin analogue only, n (%)	Human insulin and insulin analogue, n (%)
N	870	397	394
Maternal characteristics			
Maternal age, mean (SD)	29.3±5.2	29.8±5.1	29.1±5.4
Maternal age, >35 years	150 (17.2)	67 (16.9)	67 (17.0)
Multiparous	515 (59.2)	216 (54.4)	211 (53.6)
Planned pregnancy	208 (23.9)	196 (49.4)*	139 (35.3)
Type 1 diabetes	791 (90.9)	379 (95.5)	359 (91.1)
Duration of diabetes, >10 years	405 (46.6)	255 (64.2)*	188 (47.7)
Chronic hypertension	42 (4.8)	25 (6.3)	15 (3.8)
HbA1c first trimester			
≤43 mmol/mol (≤6.1%)	109 (12.5)	54 (13.6)	66 (16.8)
44–53 mmol/mol (6.2%–7.0%)	120 (13.8)	117 (29.5)*	66 (16.8)
54–67 mmol/mol (7.1%–8.3%)	134 (15.4)	86 (21.7)	94 (23.9)
>67 mmol/mol (>8.3%)	100 (11.8)	54 (13.6)	47 (11.9)
Unknown	407 (46.8)	86 (21.7)	121 (30.7)
Infant characteristics			
Birth year			
≤2002	363 (41.7)	13 (3.3)	25 (6.4)
2003–2007	357 (41.0)	126 (31.7)*	121 (30.7)*
2008–2012	150 (17.3)	258 (65.0)*	248 (62.9)*
Fetal outcome			
Live birth	855 (98.3)	380 (95.7)	380 (96.4)
Stillbirth	11 (1.3)	10 (2.5)	7 (1.8)
Spontaneous abortion	1 (0.1)	6 (1.5)*	5 (1.3)*
Termination	3 (0.3)	1 (0.3)	2 (0.5)
Male sex	456 (52.4)	200 (50.4)	208 (52.8)
Preterm delivery‡, <37 weeks	249 (29.2)	122 (32.7)	80 (21.1)*
Large for gestation‡	413 (48.7)	197 (53.4)	184 (48.6)
Death in the first week‡	6 (0.8)	1 (0.3)	0

The range of gestational age is 6–42 weeks.

*P<0.05.

†Reference group for comparison was fetuses exposed to human insulin only in the first trimester.

‡Among live birth.

HbA1c, haemoglobin A1c.

risk of major congenital anomaly (OR 0.54; 95% CI 0.29 to 1.01), but not statistically significant. The risk of major congenital anomaly was significantly lower among those exposed to both human insulin and insulin analogues (OR 0.43; 95% CI 0.21 to 0.89). Mean HbA1c level in the first trimester was significantly associated with major congenital anomaly, with OR of 2.87 (95% CI 1.28 to 6.42) for the HbA1c level above 67 mmol/mol (8.3%). Adjusting for the mean HbA1c in the first trimester and region did not change the risk estimates (table 3). The OR of congenital heart defects adjusted for region and HbA1c was 0.14 (95% CI 0.03 to 0.62) among fetuses

with exposure to insulin analogue only, and 0.47 (95% CI 0.19 to 1.19) among those with exposure to both human insulin and insulin analogue, compared with the reference group (table 3). In the analysis of adverse outcomes (major congenital anomaly or fetal death), the point estimates did not show increased risks associated with exposure to insulin analogues (either alone or with human insulin), with an adjusted OR of 0.76 (95% CI 0.44 to 1.33) for those exposed to insulin analogue only (table 3). In the sensitivity analysis, missing HbA1c levels in the first trimester (n=614) were included as a separate group in the multivariate model (online supplementary

**Table 3** Relative risk of non-chromosomal congenital anomalies by treatment of diabetes in the first trimester

	Human insulin only, N=463	Insulin analogue only, N=311	Human insulin and insulin analogue, N=273
Major congenital anomaly			
n (%)	37 (8.0)	14 (4.5)	10 (3.7)
Crude OR (95% CI)	Reference	0.54 (0.29 to 1.01)	0.43 (0.21 to 0.89)
Adjusted OR (95% CI)*	Reference	0.56 (0.29 to 1.06)	0.44 (0.22 to 0.91)
Congenital heart defects			
n (%)	21 (4.5)	2 (0.6)	6 (2.2)
Crude OR (95% CI)	Reference	0.14 (0.03 to 0.58)	0.46 (0.18 to 1.15)
Adjusted OR (95% CI)*	Reference	0.14 (0.03 to 0.62)	0.47 (0.19 to 1.19)
Adverse outcomes†			
n (%)	42 (9.1)	21 (6.8)	19 (7.0)
Crude OR (95% CI)	Reference	0.72 (0.42 to 1.24)	0.74 (0.42 to 1.31)
Adjusted OR (95% CI)*	Reference	0.76 (0.44 to 1.33)	0.77 (0.44 to 1.36)

*Adjusted for centre as a random effect and adjusted for HbA1c value in the first trimester.

†Including major congenital anomaly or fetal deaths.

HbA1c, haemoglobin A1c.

table 1). The estimated ORs of major congenital anomaly overall or congenital heart defects with insulin analogue use were similar.

Planned pregnancy was associated with a lower risk of major congenital anomalies (OR 0.73, 95% CI 0.37 to 1.46) and for congenital heart defects (OR 0.28, 95% CI 0.20 to 1.58). Adjusting for other potential confounders (maternal age, planned pregnancy, diabetes type and year of delivery) did not significantly affect the estimated ORs

of major congenital anomalies overall or congenital heart defects with insulin analogues use (data not shown).

Out of 311 fetuses exposed to insulin analogues only, 217 (69.8%) were exposed to short-acting insulin analogues only and 94 (19.2%) were exposed to a combination of long-acting analogues with short-acting insulin analogues (table 4). When analyses were restricted to fetuses exposed to insulin analogues only, the significantly lower risk of major congenital anomalies/congenital heart defects was

Table 4 ORs of non-chromosomal congenital anomalies by type of insulin analogues among fetuses exposed to insulin analogues only in the first trimester

N	Human insulin only	Insulin analogues only	
		Short-acting analogues only	Short-acting analogues and long-acting analogues
		463	217
Major congenital anomaly			
n (%)	37 (8.0)	9 (4.2)	5 (5.3)
Crude OR (95% CI)	Reference	0.49 (0.23 to 1.04)	0.64 (0.24 to 1.68)
Adjusted OR (95% CI)*	Reference	0.57 (0.26 to 1.21)	0.54 (0.20 to 1.43)
Congenital heart defects			
n (%)	21 (4.5)	1 (0.5)	1 (1.1)
Crude OR (95% CI)	Reference	0.10 (0.01 to 0.72)	0.23 (0.03 to 1.70)
Adjusted OR (95% CI)*	Reference	0.12 (0.02 to 0.92)	0.18 (0.02 to 1.35)
Adverse outcomes†			
n (%)	42 (9.1)	15 (6.9)	6 (6.4)
Crude OR (95% CI)	Reference	0.74 (0.40 to 1.36)	0.68 (0.28 to 1.64)
Adjusted OR (95% CI)*	Reference	0.94 (0.45 to 1.98)	0.53 (0.21 to 1.36)

*Adjusted for centre as a random effect and adjusted for HbA1c value in the first trimester.

†Including major congenital anomaly or fetal deaths.

HbA1c, haemoglobin A1c.

Table 5 Prevalence* of non-chromosomal congenital anomaly subgroups by treatment of diabetes in the first trimester

	Human insulin only N=870	Insulin analogues only N=397	Human insulin and insulin analogue N=394
EUROCAT subgroups	n (per 1000)	n (per 1000)	n (per 1000)
Nervous system	11 (12.6)	2 (5.0)	2 (5.1)
Ear, face and neck	1 (1.1)	0	0
Congenital heart defects	35 (40.2)	3 (7.6)	14 (35.5)
Severe congenital heart defects	2 (2.3)	0	4 (10.2)
Transposition of great vessels	0	0	2 (5.1)
Single ventricle	1 (1.1)	0	0
Ventricular septal defect	20 (23.0)	1 (2.5)	9 (22.8)
Atrial septal defect	12 (13.8)	2 (5.0)	2 (5.1)
Tetralogy of Fallot	0	0	1 (2.5)
PDA as only congenital heart defects in term infants (GA +37 weeks)	1 (1.1)	0	1 (2.5)
Respiratory	1 (1.1)	1 (2.5)	1 (2.5)
Orofacial clefts	1 (1.1)	2 (5.0)	0
Digestive system	5 (5.7)	0	2 (5.1)
Urinary	10 (11.5)	3 (7.6)	1 (2.5)
Genital	3 (3.4)	2 (5.0)	2 (5.1)
Limb	9 (10.3)	4 (10.1)	1 (2.5)
Other anomalies/syndromes	4† (4.6)	0	1‡ (2.5)

*One fetus can be counted in more than one subgroup if it has multiple malformations, but only once in total.

†Including craniosynostosis (n=2), teratogenic syndromes with malformations (n=1), genetic syndromes+microdeletions (n=1).

‡Situs inversus (n=1).

GA, gestational age; PDA, patent ductus arteriosus.

seen in fetuses exposed to short-acting insulin analogue only (OR 0.12, 95% CI 0.02 to 0.92), although the point estimates were similar between exposure to short-acting insulin analogue only and short-acting and long-acting insulin analogues (table 4).

Table 5 describes the specific congenital anomalies diagnosed in all fetuses (n=1661). Congenital heart defects were the most frequent except in those exposed to insulin analogues only. Of the three congenital heart defects cases among insulin analogues only group, two were atrial septal defect (ASD) and one ventricular septal defect (VSD). Of the 35 fetuses with congenital heart defects exposed to human insulin only, two had severe congenital heart defects and the remaining 33 cases had septal defects or persistent ductus. There were no cases with caudal regression sequence.

DISCUSSIONS

In this retrospective population-based cohort study, we found no increase in the risk of congenital anomalies in fetuses exposed to insulin analogues in the first trimester compared with those exposed to human insulin. Furthermore, a significantly lower risk of congenital heart defects was observed with exposure to insulin analogues.

In this cohort of pregnancies with diabetes, the rate of major congenital anomalies was 7.0% and about half of all major congenital anomalies were congenital heart defects. These rates are comparable with previous reports which found that the risk of congenital anomalies in the offspring of women with diabetes was between 4.6% and 9.7%,^{2-4 17} a twofold to fivefold increase in risk compared with the general population in the study region.¹⁸ Our study confirms that unplanned pregnancy² and elevated HbA1c levels in the first trimester are associated with a higher rate of congenital anomalies.^{19 20}

In Europe, insulin analogues have been increasingly used during pregnancy in the last decade,²¹ and so, it is important to assess their safety. Studies have consistently showed no increased risk of congenital anomalies overall with the use of insulin analogues in pregnancy compared with the use of human insulin.²²⁻²⁵ Among a number of retrospective and observational studies on insulin lispro, the largest was a retrospective multinational study of 533 pregnancies with pregestational diabetes,²² which reported a rate of major congenital anomalies of 5.4% (n=27, 95% CI 3.5% to 7.4%). An open-label randomised controlled trial found a 4.5% rate of congenital anomalies for insulin aspart among 322 women with type 1 diabetes,

comparable to a 6.6% rate for human insulin.²³ Most studies exploring the use of insulin glargine in pregnant women with diabetes are small, retrospective and without a reference group. A meta-analysis of eight studies on the use of insulin glargine in pregnancy showed a similar rate of congenital anomalies (8.1%) compared with the use of neutral protamine Hagedorn (NPH) (7.7%) (OR 0.97, 95% CI 0.47 to 1.99).²⁵ A multinational randomised control trial showed that the frequencies of congenital anomaly among insulin detemir or NPH were similar (insulin detemir: n=8/142, 5.6%; NPH: n=8/145, 5.5%).²⁴

Our findings confirm and extend these observations. The risk of congenital anomalies was lower in fetuses exposed to insulin analogues, in particular short-acting insulin analogue only, in the first trimester compared with those exposed to human insulin. This decrease was driven by a reduction of risk for congenital heart defects. Studies outside pregnancy have showed that the use of insulin analogues has been associated with either modestly improved glycaemic control or fewer hypoglycaemic episodes.^{10 12} Recent data have emphasised the role of glucose stability in the pathogenesis of fetal malformations, as a single day of poor glycaemic control had potential for a negative impact on organogenesis.^{26 27} It is possible that insulin analogues reduced the risk of congenital anomalies by achieving better glucose stability and consequently reduced glucose-level variation. We might speculate that patients treated with insulin analogues had less frequent very high glucose spikes or hypoglycaemic episodes that might be responsible for impaired fetal development in very early pregnancy. However, our study is lacking information on glucose variability, which may also be a confounding factor. Much remains to be learnt about the biological mechanisms that explain the observed association between insulin analogues and congenital anomalies and whether insulin analogues could reduce the risk of congenital heart defects. The rate of major congenital anomalies was 8.0% in the reference group—exposure to the human insulin only. This rate is higher compared with the rate of congenital anomalies in the recent UK National Pregnancy and Diabetes Audit: 46.2 for type 1 diabetes and 34.6 for type 2 diabetes per 1000 live births and stillbirths.²⁸ This suggests that hyperglycaemia might be more severe, more frequent or possibly due to less adherence to treatment among women with diabetes in the human insulin only group. This may have exacerbated the difference in insulin analogue and human insulin outcomes beyond purely treatment-related effects. Although we do not have a clear explanation for this higher rate, it calls for increased counselling efforts towards diabetic women of childbearing age.

In our data, pregnancies with exposure to insulin analogues had significantly higher rate of spontaneous abortion compared with exposure to human insulin, which has not been previously reported. When we combined the adverse outcomes, including fetal death and congenital anomaly, we did not see increased risks

with insulin analogues compared with human insulin. There were two cases with severe congenital heart defects among fetuses exposed to human insulin only and 0 cases of severe congenital heart defects in the insulin analogue only exposed group. These numbers are too small to interpret and formulate any conclusions. It is difficult to say which is worse scenario: the increased risk (if verified) of early-spontaneous abortion following exposure to insulin analogues or the increased risk of septal defect, that often close spontaneously, following exposure to human insulin only. There were no cases with the caudal regression sequence. We would not expect to have a case in our cohort of 1661 fetuses due to its low prevalence (1 per 50 000 births),²⁹ although caudal regression sequence is highly associated with maternal diabetes with OR >20 compared with pregnancies without diabetes.⁶

Our study established a large, retrospective cohort of pregnancies with diabetes, with detailed information on treatment of diabetes, HbA1c value and other exposure characteristics. The strength of this study is its use of multiple centres that represent geographically distinct areas where patterns of treatment of diabetes are different. Another strength of the study was the valid information on congenital anomaly gained by linking to the EUROCAT database and by reclassifying the cases with written text descriptions of a congenital anomaly based on EUROCAT subgroup definition. The EUROCAT database includes well-validated, comparable and specific information on congenital anomalies in births as well as terminations of pregnancies. In addition, the associations between some well-known risk factors (such as HbA1c and planned pregnancy) and congenital anomalies in pregnancies with diabetes were consistent with prior reports, supporting the premise that the exposures and outcomes of interest were well captured in our study.

Our results should be interpreted in the context of limitations inherent in its design and local variations. We tried to identify potential confounders of the association between treatment of diabetes and subsequent congenital anomalies, but it remains possible that residual bias is still present because of unmeasured or not perfectly measured or unknown confounders, for example, other as yet unknown differences between groups of insulin use. Our study is a retrospective cohort with information based on medical records and healthcare databases, which were not primarily designed for the current research purposes in most centres. Our data sources have incomplete data on HbA1c and pregnancy planning, which may represent a source of potential residual confounding. Fetuses of mothers with missing values of HbA1c had an increased risk of congenital anomalies/congenital heart defects in our data. In the sensitivity analysis by including 'missing HbA1c value' in the model as a separate group, the risk estimation did not show a significant difference from the analysis that excluded missing HbA1c. We were not able to adjust for preconception care (eg, screening and treatment of complications of diabetes, folic acid supplementation) which

is also known to lower the risk of congenital anomaly. In addition, it is possible that the underlying choice of insulin analogue, human insulin or the switch to insulin analogue may confound the risk estimation in the study. Information on the use of an insulin pump, which may have an important impact on the stability of glucose level control, is missing in our study. Pregnant women with renal disease might have a higher risk of congenital anomalies, but information on microvascular complications, particularly diabetic nephropathy is lacking. The cohort data were collected at centres for pregnancies with diabetes and the gestational age at referral to these centres may differ between regions. Therefore, the inclusion of spontaneous abortions may differ. The low rate of spontaneous abortion in our study may be due to the medical records not starting very early in pregnancy; therefore, early spontaneous abortion may not have been captured in our database. Spontaneous abortion was not a primary outcome of the study. Ascertainments of congenital anomalies procedures vary between registries, although we adjusted for region in the analysis. The follow-up period for inclusion of congenital anomalies varies in the EUROCAT registries. As septal defects often are diagnosed after the neonatal period depending on the local protocol, there may be a difference in the inclusion of ASD and VSD. On the other hand, regions with frequent use of echocardiography in the neonatal units may diagnose more septal defects in the neonatal period.³⁰ A further limitation of this analysis is the low number of exposed cases, and even though we applied methods appropriate for small cell counts, our ability to control confounding is limited. The absolute numbers of anomalies are small and may be a chance finding or explained by unadjusted confounders. While the biological effect varies according to each type of insulin among short-acting or long-acting insulin analogues, the low numbers of anomalies limited our ability to evaluate the risk in relation to each individual analogue. Because the database did not contain information on lifestyle factors (eg, smoking, obesity, alcohol), we were not able to examine their potential for such confoundings.

In conclusion, our results suggest that first trimester use of insulin analogues does not increase the risk of a congenital anomaly among women with diabetes compared with the use of human insulin. This is the first study that shows a significantly lower risk of congenital heart defects in relation to insulin analogue use, but caution is warranted due to the small numbers of anomalies and the as yet unknown confounders. Our study provides a further piece of evidence on the relative safety of insulin analogues with regards to congenital anomaly, in particular to congenital heart defects. The relative higher proportion of spontaneous abortion among women exposed to insulin analogues observed in our study needs to be investigated in an independent dataset as it was not the primary outcome of our study.

Author affiliations

- ¹Pharmacoepidemiology and Pharmacoeconomics, University of Groningen, Groningen, The Netherlands
- ²Division of Reproduction, Poznan University of Medical Sciences, Poznan, Poland
- ³Paediatric Department, Hospital Lillebaelt, Kolding, Denmark
- ⁴Obstetrician & Gynaecologist, Singleton Hospital, Swansea, UK
- ⁵School of Nursing, Ulster University, Northern Ireland, UK
- ⁶Queen Mary University of London, London, UK
- ⁷Faculty of Medical Science, University Medical Centre Groningen, Groningen, The Netherlands
- ⁸Department of Health Information and Research, Ministry of Health, Valletta, Malta
- ⁹School of Nursing, Swansea University, Swansea, UK
- ¹⁰Department of Medical Genetics, Poznan University of Medical Sciences, Poznan, Poland
- ¹¹Provinciaal Instituut voor Hygiene, Antwerp, Belgium
- ¹²Birth Registry Mainz Model, Children's Hospital, University Medical Center of Mainz, Mainz, Germany
- ¹³Ulster University, Northern Ireland, UK

Acknowledgements We thank Dr Thera Links (The Netherlands), Dr Bruce Wolffenbuttel (The Netherlands), Dr Loek de Heide (The Netherlands), Dr Klaas Hoogenberg (The Netherlands), Dr Annette Queißer-Wahrendorf (Germany), Dr Gabi Stolz (Germany), Rhian Thomas (UK), David Tucker (UK), Professor Anna Latos-Bielenska (Poland), Dr Pawel Gutaj (Poland), Anna Jamry-Dziurla (Poland), Zofia Kierznowska (Poland), who contributed to data collection of local cohort of pregnancies with diabetes. We thank Kacie Smith (UK) and James Densem (UK) who contributed to set up database. We thank Anne Vinkel Hansen (Denmark) who assisted with data analysis.

Contributors HW: drafting/revising the manuscript for content; study concept or design; analysis or interpretation of data; acquisition of data; statistical analysis; study supervision or coordination. EW-O: drafting/revising the manuscript for content; analysis or interpretation of data; study supervision or coordination; obtaining funding only for meetings attending. EG: reviewed all drafts of the manuscript for content; study concept or design; analysis or interpretation of data; acquisition of data; study supervision or coordination as Steering Group member; obtaining funding; attended all meetings. MM: drafting/revising the manuscript for content; study concept or design; analysis or interpretation of data; acquisition of data. ML: drafting/revising the manuscript for content; analysis of interpretation of data; acquisition of data; study supervision or coordination. JKM: drafting/revising the manuscript for content; analysis or interpretation of data; statistical analysis. MKB: drafting/revising the manuscript for content; analysis or interpretation of data; acquisition of data. MG: drafting/revising the manuscript for content; acquisition of data. HW: drafting/revising the manuscript for content; acquisition of data. SJ: drafting/revising the manuscript for content; study concept or design; analysis or interpretation of data; acquisition of data; study supervision or coordination; obtaining funding. AM-K: acquisition of data. VN and GT: drafting/revising the manuscript for content; acquisition of data. AW: drafting/revising the manuscript for content; study concept or design; analysis or interpretation of data; acquisition of data; statistical analysis; study supervision or coordination. HD: drafting/revising the manuscript for content; study concept or design; analysis of interpretation of data; statistical analysis; study supervision or coordination; obtaining funding. LTWJ-B: drafting/revising the manuscript for content; study concept or design; analysis or interpretation of data; acquisition of data; study supervision or coordination; obtaining funding; attended all meetings of EUROmedICAT.

Funding This study was part of the EUROmedICAT research project (www.euromedicat.eu) which has been supported by the European Commission under the Seventh Framework Programme Grant agreement no 260598.

Competing interests EW-O is a member of Polish Novo Nordisk advisory board, she gives lectures and leads workshops organised by Eli Lilly and Novo Nordisk in Poland.

Patient consent Not required.

Ethics approval Approval was received from an ethical standards committee in each country.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially,



and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Jensen DM, Damm P, Moelsted-Pedersen L, *et al.* Outcomes in type 1 diabetic pregnancies: a nationwide, population-based study. *Diabetes Care* 2004;27:2819–23.
- Evers IM, de Valk HW, Visser GH. Risk of complications of pregnancy in women with type 1 diabetes: nationwide prospective study in the Netherlands. *BMJ* 2004;328:915.
- Persson M, Norman M, Hanson U. Obstetric and perinatal outcomes in type 1 diabetic pregnancies: A large, population-based study. *Diabetes Care* 2009;32:2005–9.
- Macintosh MC, Fleming KM, Bailey JA, *et al.* Perinatal mortality and congenital anomalies in babies of women with type 1 or type 2 diabetes in England, Wales, and Northern Ireland: population based study. *BMJ* 2006;333:177.
- Vinceti M, Malagoli C, Rothman KJ, *et al.* Risk of birth defects associated with maternal pregestational diabetes. *Eur J Epidemiol* 2014;29:411–8.
- Garne E, Loane M, Dolk H, *et al.* Spectrum of congenital anomalies in pregnancies with pregestational diabetes. *Birth Defects Res A Clin Mol Teratol* 2012;94:134–40.
- Reece EA, Homko CJ. Why do diabetic women deliver malformed infants? *Clin Obstet Gynecol* 2000;43:32–45.
- Inkster ME, Fahey TP, Donnan PT, *et al.* Poor glycosylated haemoglobin control and adverse pregnancy outcomes in type 1 and type 2 diabetes mellitus: systematic review of observational studies. *BMC Pregnancy Childbirth* 2006;6:30.
- Wahabi HA, Alzeidan RA, Bawazeer GA, *et al.* Preconception care for diabetic women for improving maternal and fetal outcomes: a systematic review and meta-analysis. *BMC Pregnancy Childbirth* 2010;10:63.
- Horvath K, Jeitler K, Berghold A, *et al.* Long-acting insulin analogues versus NPH insulin (human isophane insulin) for type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2007;(2):CD005613.
- Hirsch IB. Insulin analogues. *N Engl J Med* 2005;352:174–83.
- Siebenhofer A, Plank J, Berghold A, *et al.* Short acting insulin analogues versus regular human insulin in patients with diabetes mellitus. *Cochrane Database Syst Rev* 2004 (2):CD003287.
- de Jong J, Garne E, Wender-Ozegowska E, *et al.* Insulin analogues in pregnancy and specific congenital anomalies: a literature review. *Diabetes Metab Res Rev* 2016;32:366–75.
- Dolk H, Jentink J, Loane M, *et al.* Does lamotrigine use in pregnancy increase orofacial cleft risk relative to other malformations? *Neurology* 2008;71:714–22.
- EUROCAT. *EUROCAT Guide 1.3 and Reference Documents, Instructions for the Registration and Surveillance of Congenital Anomalies*. Northern Ireland, UK: EUROCAT, 2015.
- Visser GH, Eilers PH, Elferink-Stinkens PM, *et al.* New Dutch reference curves for birthweight by gestational age. *Early Hum Dev* 2009;85:737–44.
- Casson IF, Clarke CA, Howard CV, *et al.* Outcomes of pregnancy in insulin dependent diabetic women: results of a five year population cohort study. *BMJ* 1997;315:275–8.
- Boyd PA, Haeusler M, Barisic I, *et al.* Paper 1: The EUROCAT network – organization and processes. *Birth Defects Res A Clin Mol Teratol* 2011;91(Suppl 1):S2–15.
- Miller JL, de Veciana M, Turan S, *et al.* First-trimester detection of fetal anomalies in pregestational diabetes using nuchal translucency, ductus venosus Doppler, and maternal glycosylated hemoglobin. *Am J Obstet Gynecol* 2013;208:385.e1–385.e8.
- Miller E, Hare JW, Cloherty JP, *et al.* Elevated maternal hemoglobin A1c in early pregnancy and major congenital anomalies in infants of diabetic mothers. *N Engl J Med* 1981;304:1331–4.
- Charlton RA, Klungsoyr K, Neville AJ, *et al.* Prescribing of Antidiabetic Medicines before, during and after Pregnancy: A Study in Seven European Regions. *PLoS One* 2016;11:e0155737.
- Wyatt JW, Frias JL, Hoyme HE, *et al.* Congenital anomaly rate in offspring of mothers with diabetes treated with insulin lispro during pregnancy. *Diabet Med* 2005;22:803–7.
- Hod M, Damm P, Kaaja R, *et al.* Fetal and perinatal outcomes in type 1 diabetes pregnancy: a randomized study comparing insulin aspart with human insulin in 322 subjects. *Am J Obstet Gynecol* 2008;198:186.e1–186.e7.
- Hod M, Mathiesen ER, Jovanović L, *et al.* A randomized trial comparing perinatal outcomes using insulin detemir or neutral protamine Hagedorn in type 1 diabetes. *J Matern Fetal Neonatal Med* 2014;27:7–13.
- Pollex E, Moretti ME, Koren G, *et al.* Safety of insulin glargine use in pregnancy: a systematic review and meta-analysis. *Ann Pharmacother* 2011;45:9–16.
- Wender-Ozegowska E, Wróblewska K, Zawiejska A, *et al.* Threshold values of maternal blood glucose in early diabetic pregnancy – prediction of fetal malformations. *Acta Obstet Gynecol Scand* 2005;84:17–25.
- de Valk HW, Visser GH. Insulin during pregnancy, labour and delivery. *Best Pract Res Clin Obstet Gynaecol* 2011;25:65–76.
- NHS UK. *National pregnancy in diabetes audit report - 2015*. London, UK: NHS UK, 2016.
- Anonymous. *Caudal regression sequence*:2016.
- Garne E, Olsen MS, Johnsen SP, *et al.* How do we define congenital heart defects for scientific studies? *Congenit Heart Dis* 2012;7:46–9.

BMJ Open

Insulin analogues use in pregnancy among women with pregestational diabetes mellitus and risk of congenital anomaly: a retrospective population-based cohort study

Hao Wang, Ewa Wender-Ozegowska, Ester Garne, Margery Morgan, Maria Loane, Joan K Morris, Marian K Bakker, Miriam Gatt, Hermien de Walle, Susan Jordan, Anna Materna-Kiryluk, Vera Nelen, Guy Thys, Awi Wiesel, Helen Dolk and Lolkje T W de Jong-van den Berg

BMJ Open 2018 8:

doi: 10.1136/bmjopen-2016-014972

Updated information and services can be found at:

<http://bmjopen.bmj.com/content/8/2/e014972>

These include:

References

This article cites 25 articles, 5 of which you can access for free at:
<http://bmjopen.bmj.com/content/8/2/e014972#ref-list-1>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Diabetes and Endocrinology](#) (456)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>